

Class 11 Maths Chapter 9. Sequences and Series

1. Sequence: Sequence is a function whose domain is a subset of natural numbers. It represents the images of $1, 2, 3, \dots, n$, as $f_1, f_2, f_3, \dots, f_n$, where $f_n = f(n)$.

2. Real Sequence: A sequence whose range is a subset of \mathbb{R} is called a real sequence.

3. Series: If $a_1, a_2, a_3, \dots, a_n$ is a sequence, then the expression $a_1 + a_2 + a_3 + \dots + a_n$ is a series.

4. Progression: A sequence whose terms follow certain rule is called a progression.

5. Finite Series: A series having finite number of terms is called finite series.

6. Infinite Series: A series having infinite number of terms is called infinite series.

Arithmetic Progression (AP)

A sequence in which the difference of two consecutive terms is constant, is called Arithmetic Progression (AP).

Properties of Arithmetic Progression

(i) If a sequence is an AP, then its n th term is a linear expression in n , i.e., its n th term is given by $An + B$, where A and B are constants and $A =$ common difference.

(ii) n th Term of an AP If a is the first term, d is the common difference and l is the last term of an AP, then

(a) n th term is given by $l = a_n = a + (n - 1)d$

(b) n th term of an AP from the last term is $a'_n = l - (n - 1)d$

(c) $a_n + a'_n = a + l$

i.e., n th term from the start + n th term from the end

= constant

= first term + last term

(d) Common difference of an AP

$$d = T_n - T_{n-1}, \forall n > 1$$

(e) $T_n = \frac{1}{2}[T_{n-k} + T_{n+k}]$, $k < n$

(iii) If a constant is added or subtracted from each term of an AP, then the resulting sequence is an AP with same common difference.

(iv) If each term of an AP is multiplied or divided by a non-zero constant k , then the resulting sequence is also an AP, with common difference kd or d/k where $d =$ common difference.

(v) If a_n, a_{n+1} and a_{n+2} are three consecutive terms of an AP, then $2a_{n+1} = a_n + a_{n+2}$.

(vi) (a) Any three terms of an AP can be taken as $a - d, a, a + d$.

(b) Any four terms of an AP can be taken as $a-3d, a- d, a + d, a + 3d$.

(c) Any five terms of an AP can be taken as $a-2d, a - d, a, a + d, a + 2d$.

(vii) Sum of n Terms of an AP

(a) Sum of n terms of AP, is given by $S_n = n/2[2a + (n - 1)d] = n/2[a + l]$

(b) A sequence is an AP, iff the sum of n terms is of the form $An^2 + Bn$, where A and B are constants. Common difference in such case will be $2A$.

(c) $T_n = S_n - S_{n-1}$

(viii) a^2, b^2 and c^2 are in AP.

$$\Leftrightarrow \frac{1}{b+c}, \frac{1}{c+a}, \frac{1}{a+b} \text{ are in AP.}$$

(ix) If a_1, a_2, \dots, a_n are the non-zero terms of an AP, then

$$\frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \frac{1}{a_3 a_4} + \dots + \frac{1}{a_{n-1} a_n} = \frac{n-1}{a_1 a_n}$$

(x) Arithmetic Mean

(a) If a, A and b are in AP, then $A = (a + b)/2$ is called the 2 arithmetic mean of a and b .

(b) If $a_1, a_2, a_3, \dots, a_n$ are n numbers, then their AM is given by,

$$d = \frac{b - a}{n + 1}$$

$$A_1 = a + d = \frac{na + b}{n + 1}$$

$$A_2 = a + 2d = \frac{(n - 1)a + 2b}{n + 1}$$

$$\dots\dots A_n = a + nd = \frac{a + nb}{n + 1}$$

(c) If $a, A_1, A_2, A_3, \dots, A_n, b$ are in AP, then $A_1, A_2, A_3, \dots, A_n$ are n arithmetic mean between a and b , where

$$d = \frac{b-a}{n+1}$$

$$A_1 = a + d = \frac{na+b}{n+1}$$

$$A_2 = a + 2d = \frac{(n-1)a+2b}{n+1}$$

$$\dots\dots A_n = a + nd = \frac{a+nb}{n+1}$$

(d) Sum of n AM's between a and b is nA

i.e., $A_1 + A_2 + A_3 + \dots + A_n = nA$

Important Results on AP

(i) If $a_p = q$ and $a_q = p$, then $a_{p+q} = 0, T_r = p + q - r$

(ii) If $pT_p = qT_q$, then $a_{p+q} = 0$

(iii) If $a_p = \frac{1}{q}$ and $a_q = \frac{1}{p}$, then $a_{pq} = 1$

(iv) If $S_p = q$ and $S_q = p$, then $S_{p+q} = -(p+q)$

(v) If $S_p = S_q$, then $S_{p+q} = 0$

Geometric Progression (GP)

A sequence in which the ratio of two consecutive terms is constant is called GP. The constant ratio is called common ratio (r).

i.e., $a_{n+1}/a_n = r, \forall n \geq 1$

Properties of Geometric Progression (GP)

(i) n th Term of a GP If a is the first term and r is the common ratio

(a) n th term of a GP from the beginning is $a_n = ar^{n-1}$

(b) n th term of a GP from the end is $a'_n = l/r^{n-1}$, l = last term

(c) If a is the first term and r is the common ratio of a GP, then the GP can be written as $a, ar, ar^2, \dots, ar^{n-1}, \dots$

(d) The n th term from the end of a finite GP consisting of m terms is ar^{m-n} , where a is the first term and r is the common ratio of the GP.

(e) $a_n a'_n = al$ i.e., n th term from the beginning \times n th term from the end = constant = first term \times last term.

(ii) If all the terms of GP be multiplied or divided by same non-zero constant, then the resulting sequence is a GP with the same common ratio.

(iii) The reciprocal terms of a given GP form a GP.

(iv) If each term of a GP be raised to same power, the resulting sequence also forms a GP.

(v) If the terms of a GP are chosen at regular intervals, then the resulting sequence is also a GP.

(vi) If $a_1, a_2, a_3, \dots, a_n$ are non-zero, non-negative term of a GP, then

(a) $GM = (a_1 a_2 a_3 \dots a_n)^{1/n}$

(b) $\log a_1, \log a_2, \log a_3, \dots, \log a_n$ are in an AP and vice-versa.

(vii) If a, b and c are three consecutive terms of a GP, then $b^2 = ac$

(viii) (a) Three terms of a GP can be taken as $a/r, a$ and ar .

(b) Four terms of a GP can be taken as $a/r^3, a/r, ar$ and ar^3 .

(c) Five terms of a GP can be taken as $a/r^2, a/r, ar$ and ar^2 .

(ix) Sum of n Terms of a GP

(a) Sum of n terms of a GP is given by

(a) Sum of n terms of a GP is given by

$$S_n = \begin{cases} \frac{\alpha(1-r^n)}{1-r}, & \text{if } |r| < 1 \\ \frac{\alpha(r^n - 1)}{r - 1}, & \text{if } |r| > 1 \\ \alpha n, & \text{if } |r| = 1 \end{cases}$$

(b) $S_n = \frac{\alpha - lr}{1-r}$ or $S_n = \frac{lr - \alpha}{r-1}, r \neq 1$

where l = last term of the GP

(c) If $|r| < 1$, then

$$S_\infty = \frac{\alpha}{1-r}$$

If $|r| \geq 1$, then it does not exist.

(x) Geometric Mean (GM)

(a) If a, G, b are in GP, then G is called the geometric mean of a and b and is given by $G = \sqrt{ab}$

(b) If $a, G_1, G_2, G_3, \dots, G_n, b$ are in GP, then $G_1, G_2, G_3, \dots, G_n$, are in GM's between a and b ,

where

$$r = \left(\frac{b}{a}\right)^{\frac{1}{n+1}}$$

$$G_1 = ar = a\left(\frac{b}{a}\right)^{\frac{1}{n+1}}$$

$$G_2 = ar^2 = a\left(\frac{b}{a}\right)^{\frac{2}{n+1}}$$

.....

$$G_n = ar^n = a\left(\frac{b}{a}\right)^{\frac{n}{n+1}}$$

(c) Product of n GM's, $G_1 \times G_2 \times G_3 \times \dots \times G_n = G^n$

Important Results on GP

(i) If $a_p = x$ and $a_q = y$, then

$$a_n = \left(\frac{x^{n-q}}{y^{n-p}}\right)^{\frac{1}{p-q}}$$

(ii) If $a_{m+n} = p$ and $a_{m-n} = q$, then

$$a_m = \sqrt{pq} \text{ and } a_n = p\left(\frac{q}{p}\right)^{\frac{m}{2n}}$$

(iii) If a, b and c are the p th, q th and r th terms of a GP, then

$$a^{q-r} \times b^{r-p} \times c^{p-q} = 1$$

(iv) n th term of $b + bb + bbb + \dots$ is

$$a_n = \frac{b}{9}(10^n - 1); b = 1, 2, \dots, 9$$

(v) n th term of $0 \cdot b + 0 \cdot bb + 0 \cdot bbb + \dots$ is

$$a_n = \frac{b}{9}(1 - 10^{-n}); b = 1, 2, \dots, 9$$

Harmonic Progression (HP)

A sequence $a_1, a_2, a_3, \dots, a_n$ of non-zero numbers is called a Harmonic Progression (HP), if the sequence $1/a_1, 1/a_2, 1/a_3, \dots, 1/a_n$ is an AP.

Properties of Harmonic Progression (HP)

(i) nth term of HP, if $a_1, a_2, a_3, \dots, a_n$ are in HP, then

(a) nth term of the HP from the beginning

$$a_n = \frac{1}{\frac{1}{a_1} + (n-1)\left(\frac{1}{a_2} - \frac{1}{a_1}\right)}$$
$$= \frac{a_1 a_2}{a_1 + (n-1)(a_1 - a_2)}$$

(b) nth term of the HP from the end

$$a'_n = \frac{1}{\frac{1}{a_n} - (n-1)\left(\frac{1}{a_2} - \frac{1}{a_1}\right)}$$

$$(c) \frac{1}{a_n} + \frac{1}{a'_n} = \frac{1}{a} + \frac{1}{a_n} = \text{constant}$$
$$= \frac{1}{\text{first term of AP}} + \frac{1}{\text{last term of AP}}$$

(d) $a_n = 1/a + (n-1)d$ are the first term and common difference of the corresponding AP.

(ii) Sum of harmonic progression does not exist.

Harmonic Mean

(i) If a, H, b are in HP, then H is called the harmonic mean of a and b i.e., $H = 2ab/(a + b)$

(ii) If $a, H_1, H_2, H_3, \dots, H_n, b$ are in HP, then

$H_1, H_2, H_3, \dots, H_n$

are n harmonic means between a and b where

$$H_1 = \frac{(n+1)ab}{a + nb},$$
$$H_2 = \frac{(n+1)ab}{2a + (n-1)b}, \dots$$

(iii) Harmonic Mean (HM) between $a_1, a_2, a_3, \dots, a_n$ is given by

$$\frac{1}{H} = \frac{1}{n} \left(\frac{1}{a_1} + \frac{1}{a_2} + \frac{1}{a_3} + \dots + \frac{1}{a_n} \right)$$

Important Results on HP

(i) If in a HP, $a_m = n$ and $a_n = m$,

$$\text{then } a_{m+n} = \frac{mn}{m+n}, a_{mn} = 1, a_p = \frac{mn}{p}$$

(ii) If in a HP, $a_p = qr$ and $a_q = pr$

$$\text{then } a_r = pq$$

(iii) If H is HM between a and b , then

$$(a) (H - 2a)(H - 2b) = H^2$$

$$(b) \frac{1}{H-a} + \frac{1}{H-b} = \frac{1}{a} + \frac{1}{b}$$

$$(c) \frac{H+a}{H-a} + \frac{H+b}{H-b} = 2$$

Properties of AM, GM and HM between Two Numbers

If A, G and H are arithmetic, geometric and harmonic means of two positive numbers a and b , then

$$(i) A = \frac{a+b}{2}, G = \sqrt{ab}, H = \frac{2ab}{a+b}$$

$$(ii) A \geq G \geq H$$

$$(iii) A, G, H \text{ are in GP and } G^2 = AH$$

(iv) If A, G, H are AM, GM and HM between three given numbers a, b and c , then the equation on having a, b and c as its root is

$$x^3 - 3Ax^2 + \frac{3G^3}{H}x - G^3 = 0,$$

$$\text{where } A = \frac{a+b+c}{3}, G = (abc)^{1/3}$$

$$\text{and } \frac{1}{H} = \left(\frac{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}{3} \right)$$

(v) If A_1, A_2 be two AM's, G_1, G_2 be two GM's and H_1, H_2 be two HM's between two numbers a and b , then

$$\frac{G_1 G_2}{H_1 H_2} = \frac{A_1 + A_2}{H_1 + H_2}$$

(vi) If A, G and H be AM, GM and HM between two numbers a and b, then

$$\frac{a^{n+1} + b^{n+1}}{a^n + b^n} = \begin{cases} A, & \text{if } n = 0 \\ G, & \text{if } n = -\frac{1}{2} \\ H, & \text{if } n = -1 \end{cases}$$

Arithmetico-Geometric Progression

A sequence in which every term is a product of a term of AP and GP is known as arithmetico-geometric progression.

The series may be written as

$$a, (a+d)r, (a+2d)r^2, (a+3d)r^3, \dots, [a+(n-1)d]r^{n-1}$$

Then,

$$S_n = \frac{a}{1-r} + \frac{dr(1-r^{n-1})}{(1-r)^2} - \frac{\{a+(n-1)d\}r^n}{1-r}, \text{ if } r \neq 1$$

$$S_n = \frac{n}{2} [2a + (n-1)d], \text{ if } r = 1$$

$$S_\infty = \frac{a}{1-r} + \frac{dr}{(1-r)^2}, \text{ if } |r| < 1$$

Sum of Arithmetico-Geometric Series

Type 1 Let $a_1 + a_2 + a_3 + \dots$ be a given series. If $a_2 - a_1, a_3 - a_2, \dots$ are in AP or GP, then a_n and S_n can be found by the method of difference.

$$\text{Let } S_n = a_1 + a_2 + a_3 + a_4 + \dots + a_n$$

$$S_n = a_1 + a_2 + a_3 + \dots + a_{n-1} + a_n$$

$$\text{So, } S_n - S_n = a_1 + (a_2 - a_1) + (a_3 - a_2) + (a_4 - a_3) + \dots + (a_n - a_{n-1}) - a_n$$

$$\Rightarrow a_n = a_1 + (a_2 - a_1) + (a_3 - a_2) + \dots + (a_n - a_{n-1})$$

$$\therefore a_n = a_1 + T_1 + T_2 + T_3 + \dots + T_{n-1}$$

where T_1, T_2, T_3, \dots are terms of new series and $S_n = \Sigma a_n$

Type 2 It is not always necessary that the series of first order of differences i.e., $a_2 - a_1, a_3 - a_2, \dots, a_n - a_{n-1}$ is always either in AP or in GP in such case.

$$\text{Let } a_1 = T_1, a_2 - a_1 = T_2, a_3 - a_2 = T_3, \dots, a_n - a_{n-1} = T_n$$

$$\text{So, } a_n = T_1 + T_2 + \dots + T_n \quad \dots\text{(i)}$$

$$a_n = T_1 + T_2 + \dots + T_{n-1} + T_n \quad \dots\text{(ii)}$$

On subtracting Eq. (i) from Eq. (ii), we get

$$T_n = T_1 + (T_2 - T_1) + (T_3 - T_2) + \dots + (T_n - T_{n-1})$$

Now, the series $(T_2 - T_1) + (T_3 - T_2) + \dots + (T_n - T_{n-1})$ is series of second order of differences and when it is either in AP or in GP, then $a_n = a_1 + \Sigma T_r$

Otherwise, in the similar way, we find series of higher order of differences and the nth term of the series.

Exponential Series

$$1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots \infty$$

The sum of the series is denoted by the number e.

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

$$e = 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

(i) e lies between 2 and 3.

(ii) e is an irrational number.

$$(iii) e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \infty$$

$$(iv) e^{-x} = 1 - \frac{x}{1!} + \frac{x^2}{2!} - \frac{x^3}{3!} + \dots \infty$$

Exponential Theorem

Let $a > 0$, then for all real value of x,

$$a^x = 1 + x(\log_e a) + \frac{x^2}{2!} (\log_e a)^2 + \frac{x^3}{3!} (\log_e a)^3 + \dots \infty$$

Logarithmic Series

$$(i) \log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \infty$$

$$\therefore \log_e(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$$

$$(ii) \log_e(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots \infty$$

$$\Rightarrow -\log_e(1-x) = x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots \infty$$

$$(iii) \log_e \left(\frac{1+x}{1-x} \right) = 2 \left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots \infty \right)$$

$$(iv) \log_e 2 = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots \infty$$

Important Points to be Remembered

- (i) In the exponential series $e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \infty$, all terms carry positive signs whereas in the logarithmic series $\log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$, the terms are alternatively positive and negative sign.
- (ii) In the exponential series, the denominators of the terms involve factorials of natural numbers. But in the logarithmic series the terms do not contain factorials.
- (iii) The exponential series is valid for all the values of x . The log series is valid when $|x| < 1$.

1. $\sum_{n=0}^{\infty} \frac{1}{n!} = e = \sum_{n=0}^{\infty} \frac{1}{(n-1)!} = \sum_{n=0}^{\infty} \frac{1}{(n-k)!} = e$
2. $\sum_{n=1}^{\infty} \frac{1}{n!} = \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \infty = e - 1$
3. $\sum_{n=2}^{\infty} \frac{1}{n!} = \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots \infty = e - 2$
4. $\sum_{n=0}^{\infty} \frac{1}{(n+1)!} = \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \infty = e - 1$
5. $\sum_{n=1}^{\infty} \frac{1}{(n+1)!} = \sum_{n=0}^{\infty} \frac{1}{(n+2)!} = \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots \infty = e - 2$
6. $\sum_{n=0}^{\infty} \frac{1}{(2n)!} = 1 + \frac{1}{2!} + \frac{1}{4!} + \frac{1}{6!} + \dots = \frac{e + e^{-1}}{2} = \sum_{n=1}^{\infty} \frac{1}{(2n-2)!}$
7. $\sum_{n=1}^{\infty} \frac{1}{(2n-1)!} = \frac{1}{1!} + \frac{1}{3!} + \frac{1}{5!} + \dots = \frac{e - e^{-1}}{2} = \sum_{n=0}^{\infty} \frac{1}{(2n+1)!}$
8. $e^{ax} = 1 + \frac{(ax)}{1!} + \frac{(ax)^2}{2!} + \frac{(ax)^3}{3!} + \dots + \frac{(ax)^n}{n!} + \dots \infty$
9. $\sum_{n=0}^{\infty} \frac{n}{n!} = e = \sum_{n=0}^{\infty} \frac{n}{n!}$
10. $\sum_{n=0}^{\infty} \frac{n^2}{n!} = 2e = \sum_{n=1}^{\infty} \frac{n^2}{n!}$
11. $\sum_{n=0}^{\infty} \frac{n^3}{n!} = 5e = \sum_{n=1}^{\infty} \frac{n^3}{n!}$
12. $\sum_{n=0}^{\infty} \frac{n^4}{n!} = 15e = \sum_{n=1}^{\infty} \frac{n^4}{n!}$
13. $\sum_{r=1}^n (a_r \pm b_r) = \sum_{r=1}^n a_r \pm \sum_{r=1}^n b_r$
14. $\sum_{r=1}^n k a_r = k \sum_{r=1}^n a_r$
15. $\sum_{r=1}^n k = k + k + \dots n \text{ times} = k_n$
16. $\sum_{r=1}^n r = 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
17. $\sum_{r=1}^n r^2 = 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$
18. $\sum_{r=1}^n r^3 = 1^3 + 2^3 + 3^3 + \dots + n^3 = \left[\frac{n(n+1)}{2} \right]^2$

24. If number of terms in AP/GP/HP are odd, then AM/GM/HM of first and last term in middle term of progression.
25. If p th, q th and r th term of geometric progression are also in geometric progression.
26. If a, b and c are in AP and also in GP, then $a=b=c$
27. If a, b and c are in AP, then xa, xb and xc are in geometric progression.

SUMMARY

Chapter 9

SEQUENCES AND SERIES

Arithmetic progression (A.P)

Standard AP $\rightarrow a, a+d, a+2d, \dots, a+(n-1)d$

$$A_n = a + (n-1)d$$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$= \frac{n}{2}(a + an)$$

Arithmetic mean A between the two numbers a and b is

$$A = \frac{a+b}{2}$$

If A_1, A_2, \dots, A_n are n A.M between the two numbers a and b,

Then $d = \frac{b-a}{n+1}$

$$A_1 = a + d = a + \frac{b-a}{n+1}$$

$$A_2 = a + 2d = a + 2 \frac{b-a}{n+1}$$

.....

$$A_n = a + nd = a + n \frac{b-a}{n+1}$$

Geometric progression (G.P)

Standard GP $\rightarrow a, ar, ar^2, \dots, ar^{n-1}$

$$A_n = ar^{n-1}$$

$$S_n = \frac{a(r^n-1)}{r-1} \text{ or } \frac{a(1-r^n)}{1-r} \quad \text{if } r \neq 1$$

$$S_\infty = \frac{a}{1-r} \quad \text{if } |r| < 1$$

If G is the GM between a and b, then $G = \sqrt{ab}$

If G_1, G_2, \dots, G_n are n G.M between the two numbers a and b ,

$$\text{then } r = \left(\frac{b}{a}\right)^{\frac{1}{n+1}}$$

$$G_1 = ar = a \left(\frac{b}{a}\right)^{\frac{1}{n+1}}$$

$$G_2 = ar^2 = a \left(\frac{b}{a}\right)^{\frac{2}{n+1}}$$

$$G_n = ar^n = a \left(\frac{b}{a} \right)^{\frac{n}{n+1}}$$

Sum to n terms of special series

$$S_n = 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

$$S_n = 1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$S_n = 1^3 + 2^3 + 3^3 + \dots + n^3 = \frac{\{n(n+1)\}^2}{4}$$

TEXT BOOK QUESTIONS

- * → Exercise 9.2 → Qns 5,7,8,11,14
- * → Exercise 9.3 → Qns 2,3,5,11,16,17,19,21,23,25
- * → Exercise 9.4 → Qns 3,4,5,6,7
- * → Misc Exercise → Qns 3,4,5,10,12,14,18,21
- ** → Exercise 9.2 → Qns 9,10,12,13,15
- ** → Exercise 9.3 → Qns 12,13,14,15,18,22,26,27,28
- ** → Exercise 9.4 → Qns 1,2,8,9,10
- ** → Misc Exercise → Qns 19,22,23,24, 25,26
- ** → Examples 4,5,6,10,13,18,21

EXTRA/ HOT QUESTIONS

1. Which term of the sequence $25, 24\frac{1}{4}, 23\frac{1}{2}, 22\frac{3}{4}, \dots$ is the first negative term. (Ans.35)
2. How many terms are identical in the two AP.
 $2,4,6, \dots$ up to 100 terms and $3,6,9, \dots$ up to 80 terms
 (Ans.33)
3. solve for x : $1+4+7+\dots+x = 590$ (Ans.x=58)
4. Find the sum of all the three digit numbers which leaves the remainder 2 when divided by 5 (Ans.98010)

5. The digits of a three digit natural number are in AP and their sum is 15

.The number obtained by reversing the digits is 396 less than the original number. Find the number.

6. If p^{th} , q^{th} , and r^{th} terms of GP are in AP. Show that p,q,r are in AP

7. If a,b,c,d are in GP, then show that $a^2 + b^2$, $b^2 + c^2$, $c^2 + d^2$ are in GP

8. Evaluate $7^{\frac{1}{2}} \times 7^{\frac{1}{4}} \times 7^{\frac{1}{8}} \dots$ to infinite terms.

9. The common ratio of a GP is $(-4/5)$ and sum to infinity is $(80/9)$. Find the first term. (Ans.7)

10. If S_1, S_2, S_3 are the sums of first n, 2n, 3n terms of a GP. Then Show that $s_1 (s_3 - s_2) = (s_2 - s_1)^2$

11. $\frac{1}{x+y}, \frac{1}{y+z}, \frac{1}{x+z}$ are in AP Show that y^2, x^2 and z^2 are in AP .

12. Find the sum of $10^3 + 11^3 + \dots + 20^3$ (Ans.42075)

13. Find the n^{th} term and the sum of n terms of the series

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots$$

14. Find the sum of n terms of $1^3 + \frac{1^3+2^3}{2} + \frac{1^3+2^3+3^3}{3} + \dots$

15. If AM and GM of roots of a quadratic equation are 8 and 5 respectively, then write the quadratic equation. (Ans. $x^2 - 16x + 25 = 0$)